



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Reports from Habana.

Assistant Surgeon Trotter reports, March 19 and 25, as follows:

Mortuary report of Habana, Cuba, for week ended March 14, 1903.

Disease.	Number of deaths.
Tuberculosis	20
Pneumonia.....	9
Bronchitis.....	3
Meningitis	10
Enteritis.....	1
Scarlet fever	2
Glanders	1

Total deaths from all causes, 101.

Week ended March 21, 1903.

Disease.	Number of deaths.
Tuberculosis	20
Pneumonia.....	6
Bronchitis.....	7
Grippe.....	1
Typhoid.....	1
Enteritis.....	6
Meningitis.....	6
Dysentery	1
Tetanus	1
Cancer.....	3

Total deaths from all causes, 113.

The SURGEON-GENERAL.

Reports from Matanzas—Whooping cough prevalent.

Passed Assistant Surgeon Von Ezdorf reports, March 17 and 24, as follows: Twenty-one deaths were reported in this city during the week ended March 14, 1903, making an annual rate of mortality of 22.75 per 1,000.

The causes of death were as follows: Tuberculosis 4, broncho-pneumonia 1, bronchitis 1, pulmonary congestion 1, entero-colitis 1, nephritis 1, endocarditis 1, heart disease; valvular, 1; other causes, 9.

No quarantinable disease was reported in this district.

Whooping cough is quite prevalent among the children, and pulmonary affections are more common in the city.

Eleven deaths were reported in this city during the week ended March 21, 1903, making an annual rate of mortality for the week of 11.91 per 1,000. The chief causes of death reported were as follows: Typhoid fever, 1; tuberculosis, 1; malarial cachexia, 1; septicæmia, 1; enteritis, acute, 1; Bright's disease, 1; heart disease, 2; other causes, 3. No quarantinable disease was reported in this district.

The SURGEON-GENERAL.